

SHARP Health Plan

Adolescents (11 to 17 years)¹

Preventive Care Checklist

Discuss your checklist with your doctor to develop your own personalized preventive care plan. Your doctor will help you determine which tests and health screenings are best for you based on your age, gender, health status and family history.

Exams		
Yearly Well Visit with Your Adolescent's Doctor (schedule at least 2 months in advance)	11 – 17	<input type="radio"/>
Vaccines ²		
COVID-19*	11 and older	<input type="radio"/>
Flu Shot (every fall)	11 and older	<input type="radio"/>
Human Papillomavirus (HPV) ³	11 – 12	<input type="radio"/>
Meningococcal B ⁴	11 – 12	<input type="radio"/>
Meningococcal Conjugate (MCV4) ³	11 – 12	<input type="radio"/>
Tetanus, Diphtheria, Pertussis (Tdap) ³	11 – 12	<input type="radio"/>
Screenings		
Body Mass Index (BMI)	11 and older	<input type="radio"/>
Cardiac Arrest	11 – 17	<input type="radio"/>
Depression and Suicide Risk	12 and older	<input type="radio"/>
Hepatitis B	11 and older	<input type="radio"/>
HIV ⁴	15 and older	<input type="radio"/>
Obesity	11 and older	<input type="radio"/>
Sexually Transmitted Infections (STIs) ⁵	Sexually active adolescents who are at increased risk for sexually transmitted infections (STIs)	<input type="radio"/>
Skin Cancer	11 and older	<input type="radio"/>
Tobacco, Drug and Alcohol Use Counseling	11 and older	<input type="radio"/>

* Some COVID-19 vaccines are authorized under EUA (emergency use authorization). ¹The information in these checklists includes recommendations adapted from the following sources as of January 2023 and is subject to change: Preventive services with a rating of A or B from the U.S. Preventive Services Task Force; immunization for children, adolescents and adults recommended by the Centers for Disease Control and Prevention; and preventive care screenings for infants, children, adolescents and women supported by the Health Resources and Services Administration. ² Timing of vaccines may vary based on PCP's recommendation. ³ Catch-up doses can be given after age 12 throughout adolescence. ⁴ Based on risk factors, and as advised by your doctor. ⁵ Counseling for all sexually active adolescents recommended.